

Pyridostigmine Br Oral Solution, USP, 60 mg/5 mL

S.No.	Category	Question	Answer	
	Clinical Particulars			
1	Use/Indication	What is the product indicated for?	Pyridostigmine bromide oral solution is useful in the treatment of myasthenia gravis.	
2	Dosage	What is the recommended dosage?	The recommended dose of Pyridostigmine Bromide Oral Solution are: raspberry-flavoured, containing 60 mg pyridostigmine bromide per teaspoonful (5 mL). This form permits accurate dosage adjustment for children and "brittle" myasthenic patients who require fractions of 60 mg doses. It is more easily swallowed, especially in the morning, by patients with bulbar involvement. The average dose is ten 5 mL teaspoonfuls daily, spaced to provide maximum relief when maximum strength is needed. In severe cases as many as 25 teaspoonfuls a day may be required, while in mild cases one to six teaspoonfuls a day may suffice.	
3	Administration	How do I take it?	The average dose is ten 5 mL teaspoonfuls daily, In severe cases as many as 25 teaspoonfuls a day may be required, while in mild cases one to six teaspoonfuls a day may suffice.	
4	Formulation	Can tablet be crushed?	NO	
5	Administration	What do I do if I miss a dose?	If you miss a dose of Pyridostigmine bromide oral solution continue your prescribed course of therapy, and contact your physician immediately.	
6	Administration	Use in Pediatric Population	Safety and effectiveness in pediatric patients have not been established.	
7	Administration	Use in Geriatric Population	There were too few elderly patients (n=18) in partial-onset seizure trials to assess if they respond differently from younger patients. No age-based dose adjustment for lacosamide is required, but elderly patients should start at a lower dose due to increased risks of decreased liver and kidney function, cardiac issues, and polypharmacy	



acctylcholine by cholimesterase and thereby permits free transmission of nerve impulses across the neuromuscul junction. Pyridostigmine is an analog of neostigmin (Prostigmin), but differs from it in certain clinical junction. Pyridostigmine is an analog of neostigmin (Prostigmin), but differs from it in certain clinical significant respects; for example, pyridostigmine in characterized by a longer duration of action and fewe gastrointestinal side effects. None 10 Lactation Use in Lactation Use in Lactation Use in Lactation The safety of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore use of pyridostigmine bromide in women who may become pregnant requires weighing the drug's potential benefit against its possible hazards to mother and child. 11 Pregnancy Use in Pregnancy The safety of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide in women who may become pregnant requires weighing the drug's potential benefits against its possible hazards to mother and child. 12 Precautions Is there any interaction and alcohol? 13 Interaction Is there any interaction with food? 14 Side effects What are the common side effects? None 15 Lactation 16 Lactation Is there any interaction with food? 17 Lactation 18 Lactation Interaction Is there any interaction with food? 19 Lactation and alcohol? 10 Lactation 10 Lactation 11 Lactation Is there any interaction with food? 11 Lactation Is there any interaction with food? 12 Precautions Is there any interaction with food? 13 Interaction Is there any interaction with food? 14 Side effects None 15 Lactation None 16 Lactation 17 Lactation L	8	Mechanism	Mechanism of Action	Pyridostigmine bromide inhibits the destruction of
junction. Pyridostigmine is an analog of neostigmin (Prostigmin), but differs from it in certain clinically significant respects; for example, pyridostigmine is an analog of neostigmine (Prostigmin), but differs from it in certain clinically significant respects; for example, pyridostigmine is an analog of neostigmine (Prostigmine) in the pyridostigmine pyridostigmine pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore use of pyridostigmine bromide in women who may become pregnant requires weighing the drug's potential benefit against its possible hazards to mother and child. The safety of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide in women who may become pregnant requires weighing the drug's potential benefits against its possible hazards to mother and child. None Is there any interaction between medication and alcohol? What are the common side effects? Side Effects of Pyridostigmine Bromide 1. Muscarinic Side Effects: Nausea Vomiting Diarrhea Abdominal cramps Increased peristalsis Niosis (pupil constriction) Diaphoresis (excessive sweating) 2. Nicotinic Side Effects: Muscle cramps Fasciculation (muscle twitching) Muscle weakness 3. Counteraction:				acetylcholine by cholinesterase and thereby permits freer
Prostigmin, but differs from it in certain clinically significant respects; for example, pyridostigmine in characterized by a longer duration of action and fewe gastrointestinal side effects. Program Share				transmission of nerve impulses across the neuromuscular
significant respects; for example, pyridostigmine in characterized by a longer duration of action and fewe gastrointestinal side effects. None Use in Lactation Use in Lactation Use in Lactation Use in Pregnancy Use in Pregnancy Use in Pregnancy Use in Pregnancy Is there any interaction between medication and alcohol? Interaction Is there any interaction with food? What are the common side effects? None Side effects What are the common side effects? None Side Effects of Pyridostigmine Bromide None Side Effects None Side Effects None Side Effects None Side Effects None None None None None None None None None 13 Interaction Is there any interaction with food? None				junction. Pyridostigmine is an analog of neostigmine
characterized by a longer duration of action and fewe gastrointestinal side effects. None 10 Lactation Use in Lactation Use in Lactation The safety of pyridostigmine bromide during pregnancy of lactation in humans has not been established. Therefore use of pyridostigmine bromide in women who may become pregnant requires weighing the drug's potential benefit against its possible hazards to mother and child. 11 Pregnancy Use in Pregnancy The safety of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Th				
gastrointestinal side effects.				
10				
The safety of pyridostigmine bromide during pregnancy of lactation in humans has not been established. Therefore use of pyridostigmine bromide in women who may become pregnant requires weighing the drug's potential benefit against its possible hazards to mother and child. The safety of pyridostigmine bromide in women who may become pregnant requires weighing the drug's potential benefits against its possible hazards to mother and child. Therefore, use of pyridostigmine bromide in women who may become pregnant requires weighing the drug's potential benefits against its possible hazards to mother and child. None 12		177	D1 1 D W '	8
lactation in humans has not been established. Therefore use of pyridostigmine bromide in women who may become pregnant requires weighing the drug's potential benefit against its possible hazards to mother and child. 11 Pregnancy Use in Pregnancy The safety of pyridostigmine bromide during pregnancy of a lactation in humans has not been established. Therefore, use of pyridostigmine bromide during pregnancy of a lactation in humans has not been subsished. Therefore, use of pyridostigmine bromide in women who may become pregnant requires weighing the drug's potential benefits against its possible hazards to mother and child. None 12 Precautions Is there any interaction between medication and alcohol? None 13 Interaction Is there any interaction with food? None 14 Side effects What are the common side effects? 1. Muscarinic Side Effects: Nausea Vomiting Diarrhea Abdominal cramps Increased peristalsis Increased perionchial secretions Miosis (pupil constriction) Diaphoresis (excessive sweating) 2. Nicotinic Side Effects: Muscle cramps Fasciculation (muscle twitching) Muscle weakness 3. Counteraction:				
use of pyridostigmine bromide in women who may become pregnant requires weighing the drug's potential benefit against its possible hazards to mother and child. 11 Pregnancy Use in Pregnancy Use in Pregnancy It is pregnancy It is there any interaction between medication and alcohol? It is there any interaction with food? It is a there any against its possible hazards to mother and child. None Side Effects None It is a feet of Pyridostigmine bromide in women who may become pregnant requires weighing the drug's potential benefits against its possible hazards to mother and child. None It is a feet of Pyridostigmine bromide in women who may become pregnant requires weighing the drug's potential benefits against its possible hazards to mother and child. It is a feet of Pyridostigmine bromide in women who may be	10	Lactation	Use in Lactation	
pregnant requires weighing the drug's potential benefit against its possible hazards to mother and child. The safety of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide in women who may become pregnant requires weighing the drug's potential benefits against its possible hazards to mother and child. Precautions Is there any interaction between medication and alcohol? Is there any interaction with food? Side effects What are the common side effects? What are the common side effects? I. Muscarinic Side Effects: Nausea Vomiting Diarrhea Abdominal cramps Increased peristalsis Increased peristalsis Increased peristalsis Increased peristalsis Increased pronchial secretions Miosis (pupil constriction) Diaphoresis (excessive sweating) Z. Nicotinic Side Effects: Muscle cramps Fasciculation (muscle twitching) Muscle weakness 3. Counteraction:				· · · · · · · · · · · · · · · · · · ·
against its possible hazards to mother and child. The safety of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide in women who may become pregnant requires weighing the drug's potential benefits against its possible hazards to mother and child. None Is there any interaction between medication and alcohol? Is there any interaction with food? Side effects What are the common side effects? I. Muscarinic Side Effects: Nausea Vomiting Diarrhea Abdominal cramps Increased pristalsis Increased salivation Increased pronchial secretions Miosis (pupil constriction) Diaphoresis (excessive sweating) 2. Nicotinic Side Effects: Muscle cramps Fasciculation (muscle twitching) Muscle weakness 3. Counteraction:				
The safety of pyridostigmine bromide during pregnancy or lactation in humans has not been established. Therefore, use of pyridostigmine bromide in women who may become pregnant requires weighing the drug's potential benefits against its possible hazards to mother and child. Precautions Is there any interaction between medication and alcohol? Is there any interaction with food? None None None Side effects What are the common side effects? I. Muscarinic Side Effects: Nausea None None None None None 14 Side effects None None None 15 Interaction with food? 16 Interaction with food? None None None None None None 17 Interaction with food? None None None None None None None 18 Effects of Pyridostigmine Bromide 19 Interaction Bromide None 10 Interaction What are the common side effects: Nausea None None None 10 Interaction What are the common side effects: Nausea None None None None None 10 Interaction What are the common side effects: None None None None None None 10 Interaction What are the common side effects: None				
lactation in humans has not been established. Therefore, use of pyridostigmine bromide in women who may become pregnant requires weighing the drug's potential benefits against its possible hazards to mother and child. 12 Precautions Is there any interaction between medication and alcohol? 13 Interaction Is there any interaction with food? What are the common side effects? Side Effects of Pyridostigmine Bromide 1. Muscarinic Side Effects: Nausea Vomiting Diarrhea Abdominal cramps Increased peristalsis Increased pristalsis Increased salivation Increased bronchial secretions Miosis (pupil constriction) Diaphoresis (excessive sweating) 2. Nicotinic Side Effects: Muscle cramps Fasciculation (muscle twitching) Muscle weakness 3. Counteraction:	11	Pregnancy	Use in Pregnancy	
pregnant requires weighing the drug's potential benefits against its possible hazards to mother and child. Precautions Is there any interaction between medication and alcohol? Is there any interaction with food? Is there any interaction with food? Side effects What are the common side effects? I. Muscarinic Side Effects: Nausea Vomiting Diarrhea Abdominal cramps Increased peristalsis Increased peristalsis Increased pronchial secretions Miosis (pupil constriction) Diaphoresis (excessive sweating) 2. Nicotinic Side Effects: Muscle cramps Fasciculation (muscle twitching) Muscle weakness 3. Counteraction:				
against its possible hazards to mother and child. Precautions				use of pyridostigmine bromide in women who may become
12 Precautions Is there any interaction between medication and alcohol? 13 Interaction Is there any interaction with food? 14 Side effects What are the common side effects? 15 Interaction Side Effects of Pyridostigmine Bromide 16 Interaction Is there any interaction with food? 14 Side effects What are the common side effects? 15 Interaction Interaction 16 Interaction Interaction 17 Interaction Interaction 18 Interaction Interaction 10 Interaction Interaction 11 Interaction Interaction Interaction 12 Interaction Interaction Interaction 13 Interaction Interaction Interaction Interaction 14 Interaction Interaction Interaction Interaction Interaction 15 Interaction Inte				
interaction between medication and alcohol? Is there any interaction with food? 14 Side effects What are the common side effects? Side Effects of Pyridostigmine Bromide 1. Muscarinic Side Effects: Nausea Vomiting Diarrhea Abdominal cramps Increased peristalsis Increased pristalsis Increased bronchial secretions Miosis (pupil constriction) Diaphoresis (excessive sweating) 2. Nicotinic Side Effects: Muscle cramps Fasciculation (muscle twitching) Muscle weakness 3. Counteraction:				· ·
medication and alcohol? Is there any interaction with food? What are the common side effects? None 14 Side effects What are the common side effects? Nausca Vomiting Diarrhea Abdominal cramps Increased peristalsis Increased salivation Increased bronchial secretions Miosis (pupil constriction) Diaphoresis (excessive sweating) 2. Nicotinic Side Effects: Muscle cramps Fasciculation (muscle twitching) Muscle weakness 3. Counteraction:	12	Precautions		None
alcohol? Is there any interaction with food? What are the common side effects? None I. Muscarinic Side Effects: Nausea Vomiting Diarrhea Abdominal cramps Increased peristalsis Increased salivation Increased bronchial secretions Miosis (pupil constriction) Diaphoresis (excessive sweating) 2. Nicotinic Side Effects: Muscle cramps Fasciculation (muscle twitching) Muscle weakness 3. Counteraction:				
Is there any interaction with food? None What are the common side effects? Nausea Nomiting Diarrhea Abdominal cramps Increased peristalsis Increased salivation Increased sulvation Increased sulvation Diaphoresis (excessive sweating) None None None Side Effects of Pyridostigmine Bromide Nausea Nomiting Diarrhea Abdominal cramps Increased peristalsis Increased print pronchial secretions Miosis (pupil constriction) Diaphoresis (excessive sweating) None None None None None Nuscae Nuscae Nuscae Nomiting None None Nuscae None None Nuscae Nuscae None None None None None Nuscae None N				
interaction with food? What are the common side effects? I. Muscarinic Side Effects: Nausea Vomiting Diarrhea Abdominal cramps Increased peristalsis Increased bronchial secretions Miosis (pupil constriction) Diaphoresis (excessive sweating) Increased Effects: Muscle cramps Fasciculation (muscle twitching) Muscle weakness Counteraction:	12	Interaction		None
Side effects What are the common side effects? 1. Muscarinic Side Effects: Nausea Vomiting Diarrhea Abdominal cramps Increased peristalsis Increased peristalsis Increased bronchial secretions Miosis (pupil constriction) Diaphoresis (excessive sweating) 2. Nicotinic Side Effects: Muscle cramps Fasciculation (muscle twitching) Muscle weakness 3. Counteraction:	13	Interaction		None
common side effects? 1. Muscarinic Side Effects: Nausea Vomiting Diarrhea Abdominal cramps Increased peristalsis Increased salivation Increased bronchial secretions Miosis (pupil constriction) Diaphoresis (excessive sweating) 2. Nicotinic Side Effects: Muscle cramps Fasciculation (muscle twitching) Muscle weakness 3. Counteraction:	14	Side effects		Side Effects of Pyridostigmine Bromide
 Nausea Vomiting Diarrhea Abdominal cramps Increased peristalsis Increased salivation Increased bronchial secretions Miosis (pupil constriction) Diaphoresis (excessive sweating) 2. Nicotinic Side Effects: Muscle cramps Fasciculation (muscle twitching) Muscle weakness 3. Counteraction:			common side effects?	, c
 Vomiting Diarrhea Abdominal cramps Increased peristalsis Increased salivation Increased bronchial secretions Miosis (pupil constriction) Diaphoresis (excessive sweating) 2. Nicotinic Side Effects: Muscle cramps Fasciculation (muscle twitching) Muscle weakness 3. Counteraction:				1. Muscarinic Side Effects:
 Vomiting Diarrhea Abdominal cramps Increased peristalsis Increased salivation Increased bronchial secretions Miosis (pupil constriction) Diaphoresis (excessive sweating) 2. Nicotinic Side Effects: Muscle cramps Fasciculation (muscle twitching) Muscle weakness 3. Counteraction:				Nausea
 Abdominal cramps Increased peristalsis Increased salivation Increased bronchial secretions Miosis (pupil constriction) Diaphoresis (excessive sweating) 2. Nicotinic Side Effects: Muscle cramps Fasciculation (muscle twitching) Muscle weakness 3. Counteraction:				
 Increased peristalsis Increased salivation Increased bronchial secretions Miosis (pupil constriction) Diaphoresis (excessive sweating) 2. Nicotinic Side Effects: Muscle cramps Fasciculation (muscle twitching) Muscle weakness 3. Counteraction:				Diarrhea
 Increased salivation Increased bronchial secretions Miosis (pupil constriction) Diaphoresis (excessive sweating) 2. Nicotinic Side Effects: Muscle cramps Fasciculation (muscle twitching) Muscle weakness 3. Counteraction:				Abdominal cramps
 Increased bronchial secretions Miosis (pupil constriction) Diaphoresis (excessive sweating) 2. Nicotinic Side Effects: Muscle cramps Fasciculation (muscle twitching) Muscle weakness 3. Counteraction:				
 Miosis (pupil constriction) Diaphoresis (excessive sweating) Nicotinic Side Effects: Muscle cramps Fasciculation (muscle twitching) Muscle weakness Counteraction: 				
 Diaphoresis (excessive sweating) 2. Nicotinic Side Effects: Muscle cramps Fasciculation (muscle twitching) Muscle weakness 3. Counteraction: 				
2. Nicotinic Side Effects:				
 Muscle cramps Fasciculation (muscle twitching) Muscle weakness 3. Counteraction:				Diaphoresis (excessive sweating)
 Fasciculation (muscle twitching) Muscle weakness 3. Counteraction:				2. Nicotinic Side Effects:
 Fasciculation (muscle twitching) Muscle weakness 3. Counteraction:				• Musala aramps
Muscle weakness 3. Counteraction:				
3. Counteraction:				
				- Masele Weakiess
Muscarinic side effects can typically be				3. Counteraction:
				Muscarinic side effects can typically be
				counteracted with atropine; however, this approach



			AN MISTE COMITANT
			carries some risks.
			4. Skin Reactions:
			Possible skin rash in some patients, usually subsiding after discontinuation of the medication.
			Note: Side effects are primarily related to overdosage.
15	Storage	What are the storage conditions?	Store at 25°C (77°F); excursions permitted to 15° to 30°C (59°to 86°F). [see USP Controlled Room Temperature]. Dispense in tight, light-resistant containers as defined by the USP.
16	Dispensing	How to Dispense?	As prescribed by the Physician
17	Contraindication	What are the contraindications of (medication).	, , ,
		Pharmace	eutical Particulars
18	Pharmaceutical Form	How is it supplied?	Pyridostigmine Bromide Oral Solution USP, 60 mg pyridostigmine bromide per teaspoonful (5 mL) is available as a clear, raspberry flavored solution and 0.23% w/w alcohol. It is supplied as follows: Bottles of 16 fluid ounces, (1 pint, 473 mL) NDC 72205-098-72
19	Ingredients	Active and Inactive Ingredients	Active PYRIDOSTIGMINE BROMIDE Inactive: ALCOHOL, GLYCERIN, LACTIC ACID, SODIUM BENZOATE, SORBITOL, SUCROSE, WATER
20	Coating	What is the type of coating?	None
		A	Allergens
21	Ingredients	Is it Vegetarian?	yes
22	Ingredients	Does it contain Gluten?	No
23	Ingredients	Does it contain Dairy Products?	No
24	Ingredients	Does it contain Casein	No
25	Ingredients	Does it contain Whey?	No
26	Ingredients	Does it contain corn?	No
		•	



			AN MINE COMPANY
27	Ingredients	Does it contain rye?	No
28	Ingredients	Does it contain sugar?	No
29	Ingredients	Does it contain Oats?	No
30	Ingredients	Does it contain wheat?	No
31	Ingredients	Does it contain spelt?	No
32	Ingredients	Does it contain barley?	No
33	Ingredients	Does it contain rennet?	No
34	Ingredients	Does it contain starch?	No
35	Ingredients	Does it contain Iodine?	No
36	Ingredients	Does it contain latex?	No
37	Ingredients	Does it contain alcohol?	Yes
38	Ingredients	Does it contain dyes?	No
39	Ingredients	Does it contain flavor?	No
40	Ingredients	Does it contain Lactose?	No
41	Ingredients	Does it contain Nuts?	No
42	Ingredients	Does it contain Preservatives?	Yes
43	Ingredients	Does it contain Soy products?	No
44	Ingredients	Does it contain peanut?	No
45	Ingredients	Does it contain nickel?	No
46			
Miscell aneous	Miscellaneous	May I know the product availability?	Novadoz Pharmaceuticals products are only available through pharmacies, wholesalers, and other authorized distributors. See our ADR (authorized distributors of record) page at NovadozPharma.com to learn more about where to find our products.
47	Miscellaneous	May I know about return, refunds and reimbursement?	Contact Novadoz Pharmaceuticals Customer Service directly at 908-360-1500
48	Miscellaneous	Do you have any patient's assistance program?	Novadoz Pharmaceuticals does not offer patient assistance programs at this time. The company that produces the brand version of your product may or may not offer such a program. Please check for access & eligibility requirements with that company.



49	Miscellaneous	How do I report an adverse drug effect or reaction to Novadoz medication?	To report suspected adverse reactions, contact Novadoz Pharmaceuticals LLC at 1-855-668-2369 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.
50	Miscellaneous	Why does my pharmacy that used to fill your generic formulation of a particular medicine, no longer fills my prescription with Novadoz formulation?	prescription is not a Novadoz Pharmaceuticals product. You may refer to NovadozPharma.com ADR (authorized distributor of record) page to learn where to
51	Miscellaneous	Manufacturer and Distributor	Manufactured by: MSN Pharmaceuticals Inc. Piscataway, NJ 08854 Distributed by: Novadoz Pharmaceuticals LLC Piscataway, NJ 08854-3714